Asthma Treatment Plan – Student

(This asthma action plan meets NJ Law N.J.S.A. 18A:40-12.8) **(Physician's Orders)**

The Pediatric/Adult Asthma Coalition of New Jersey
"Your Pathway to Asthma Control"
PACNJ approved Plan available at www.pacnj.org

Sponsored by American Lung Association. NJ Health New Jersey Department of Health

(Please Print)

Name	Date of Birth	Effective Date

Doctor	Parent/Guardian (if applicable)	Emergency Contact
Phone	Phone	Phone

HEALTHY (Green Zone)

Take daily control medicine(s). Some inhalers may be more effective with a "spacer" – use if directed.

You have *all* of these:
- Breathing is good
- No cough or wheeze
- Sleep through the night
- Can work, exercise, and play

And/or Peak flow above __________

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
☐ Advair® HFA ☐ 45, ☐ 115, ☐ 230	2 puffs twice a day
☐ Aerospan™	☐ 1 ☐ 2 puffs twice a day
☐ Alvesco® ☐ 80, ☐ 160	☐ 1 ☐ 2 puffs twice a day
☐ Dulera® ☐ 100, ☐ 200	2 puffs twice a day
☐ Flovent® ☐ 44, ☐ 110, ☐ 220	2 puffs twice a day
☐ Qvar® ☐ 40, ☐ 80	☐ 1 ☐ 2 puffs twice a day
☐ Symbicort® ☐ 80, ☐ 160	☐ 1 ☐ 2 puffs twice a day
☐ Advair Diskus® ☐ 100, ☐ 250, ☐ 500	1 inhalation twice a day
☐ Asmanex® Twisthaler® ☐ 110, ☐ 220	☐ 1 ☐ 2 inhalations ☐ once ☐ twice a day
☐ Flovent® Diskus® ☐ 50 ☐ 100 ☐ 250	1 inhalation twice a day
☐ Pulmicort Flexhaler® ☐ 90, ☐ 180	☐ 1 ☐ 2 inhalations ☐ once ☐ twice a day
☐ Pulmicort Respules® (Budesonide) ☐ 0.25, ☐ 0.5, ☐ 1.0	1 unit nebulized ☐ once ☐ twice a day
☐ Singulair® (Montelukast) ☐ 4, ☐ 5, ☐ 10 mg	1 tablet daily
☐ Other	
☐ None	

Remember to rinse your mouth after taking inhaled medicine.

If exercise triggers your asthma, take______________________ ____ puff(s) ____minutes before exercise.

CAUTION (Yellow Zone)

Continue daily control medicine(s) and ADD quick-relief medicine(s).

You have *any* of these:
- Cough
- Mild wheeze
- Tight chest
- Coughing at night
- Other:__________

If quick-relief medicine does not help within 15-20 minutes or has been used more than 2 times and symptoms persist, call your doctor or go to the emergency room.

And/or Peak flow from______ to______

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
☐ Albuterol MDI (Pro-air® or Proventil® or Ventolin®)	2 puffs every 4 hours as needed
☐ Xopenex®	2 puffs every 4 hours as needed
☐ Albuterol ☐ 1.25, ☐ 2.5 mg	1 unit nebulized every 4 hours as needed
☐ Duoneb®	1 unit nebulized every 4 hours as needed
☐ Xopenex® (Levalbuterol) ☐ 0.31, ☐ 0.63, ☐ 1.25 mg	1 unit nebulized every 4 hours as needed
☐ Combivent Respimat®	1 inhalation 4 times a day
☐ Increase the dose of, or add:	
☐ Other	

- **If quick-relief medicine is needed more than 2 times a week, except before exercise, then call your doctor.**

EMERGENCY (Red Zone)

Take these medicines NOW and CALL 911.
Asthma can be a life-threatening illness. Do not wait!

Your asthma is getting worse fast:
- Quick-relief medicine did not help within 15-20 minutes
- Breathing is hard or fast
- Nose opens wide • Ribs show
- Trouble walking and talking
- Lips blue • Fingernails blue
- Other:______________

And/or Peak flow below ________

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
☐ Albuterol MDI (Pro-air® or Proventil® or Ventolin®)	4 puffs every 20 minutes
☐ Xopenex®	4 puffs every 20 minutes
☐ Albuterol ☐ 1.25, ☐ 2.5 mg	1 unit nebulized every 20 minutes
☐ Duoneb®	1 unit nebulized every 20 minutes
☐ Xopenex® (Levalbuterol) ☐ 0.31, ☐ 0.63, ☐ 1.25 mg	1 unit nebulized every 20 minutes
☐ Combivent Respimat®	1 inhalation 4 times a day
☐ Other	

Triggers

Check all items that trigger patient's asthma:

- ❑ Colds/flu
- ❑ Exercise
- ❑ Allergens
  - ○ Dust Mites, dust, stuffed animals, carpet
  - ○ Pollen - trees, grass, weeds
  - ○ Mold
  - ○ Pets - animal dander
  - ○ Pests - rodents, cockroaches
- ❑ Odors (Irritants)
  - ○ Cigarette smoke & second hand smoke
  - ○ Perfumes, cleaning products, scented products
  - ○ Smoke from burning wood, inside or outside
- ❑ Weather
  - ○ Sudden temperature change
  - ○ Extreme weather - hot and cold
  - ○ Ozone alert days
- ❑ Foods:
  - ○ __________
  - ○ __________
  - ○ __________
- ❑ Other:
  - ○ __________
  - ○ __________
  - ○ __________

This asthma treatment plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs.

Disclaimers: The use of this Website/PACNJ Asthma Treatment Plan and its content is at your own risk. The content is provided on an "as is" basis. The American Lung Association of the Mid-Atlantic (ALAM-A), the Pediatric/Adult Asthma Coalition of New Jersey and all affiliates disclaim all warranties, express or implied, statutory or otherwise, including but not limited to the implied warranties of merchantability [illegible] ALAM-A makes no representations or warranties about the accuracy, reliability, completeness, currency, or timeliness of the content. [illegible]

The Pediatric/Adult Asthma Coalition of New Jersey, sponsored by the American Lung Association in New Jersey. This publication was supported by a grant from the New Jersey Department of Health and Senior Services, with funds provided by the U.S. Centers for Disease Control and Prevention [illegible] Although this document has been funded wholly or in part by the United States Environmental Protection Agency [illegible] Information in this publication is not intended to diagnose health problems or take the place of medical advice. [illegible]

REVISED AUGUST 2014

Permission to Self-administer Medication:

☐ This student is capable and has been instructed in the proper method of self-administering of the non-nebulized inhaled medications named above in accordance with NJ Law.

☐ This student is not approved to self-medicate.

PHYSICIAN/APN/PA SIGNATURE______________________ DATE______

Physician's Orders

PARENT/GUARDIAN SIGNATURE______________________

PHYSICIAN STAMP

Save

Print

Print Medicine Only